

Citation: Núñez FJ, Santalla A, Carrasquila I, Asian JA, Reina JI, Suarez-Arrones LJ (2018) The effects of unilateral and bilateral eccentric overload training on hypertrophy, muscle power and COD performance, and its determinants, in team sport players. PLoS ONE 13(3): e0193841. https://doi.org/10.1371/journal.pone.0193841

Editor: Jaime Sampaio, Universidade de Tras-os-Montes e Alto Douro, PORTUGAL

Received: November 17, 2017

Accepted: February 19, 2018

Published: March 28, 2018

Copyright: © 2018 Núñez et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: The authors received no specific funding for this work

Competing interests: The authors have declared that no competing interests exist

RESEARCH ARTICLE

The effects of unilateral and bilateral eccentric overload training on hypertrophy, muscle power and COD performance, and its determinants, in team sport players

Francisco Javier Núñez^{1®}*, Alfredo Santalla^{1,2,3®}, Irene Carrasquila¹, Jose Antonio Asian^{1,2}, Jose Ignacio Reina⁴, Luis Jesús Suarez-Arrones^{1,2®}

Department of Sports and Computing, Sport Faculty, Pablo de Olavide University of Sevilla, Sevilla, Spain,
Master de Futbol, Pablo de Olavide University of Sevilla, Sevilla, Spain,
Research Institute of Hospiltal 12 de Octubre (i+ 12), Madrid, Spain,
Dos Hermanas Radiological Center, Sevilla, Spain

 $\ensuremath{\mathfrak{O}}$ These authors contributed equally to this work.

* fjnunsan@upo.es

Abstract

The study aimed to compare the chronic eccentric-overload training effects of unilateral (lateral lunge) vs bilateral (half-squat) using an inertial device, on hypertrophy and physical performance. Twenty-seven young team sports male players performed a 4 sets of 7 repetitions of inertial eccentric overload training, biweekly for 6 weeks, distributed in unilateral lunge group (UG: age: 22.8 ± 2.9 years; body mass: 75.3 ± 8.8 kg; height: 177.3 ± 3.7 cm) and bilateral squat group (BG: age: 22.6 ± 2.7 years; body mass: 79.5 ± 12.8 kg; height: 164.2 ± 7 cm). Lower limb muscle volume, counter movement jump (CMJ), power with both (POWER), dominant (POWERd) and no-dominant leg (POWERnd), change of direction turn of 90° with dominant (COD90d) and no-dominant leg (COD90nd) and 180° (COD180d and COD180nd), and 10m sprint time (T-10m) were measured pre and post-intervention. The UG obtained an increase of adductor major (+11.1%) and vastus medialis (+12.6%) higher than BG. The BG obtained an increase of vastus lateralis (+9.9%) and lateral gastrocnemius (+9.1%) higher than UG. Both groups improved CMJ, POWER, POWERd, POWERnd, COD90 and DEC-COD90, without changes in T-10m. The UG decrease DEC-COD90nd (-21.1%) and BG increase POWER (+38.6%) substantially more than the other group. Six-weeks of unilateral / bilateral EO training induce substantial improvements in lower limbs muscle volume and functional performance, although unilateral training seems to be more effective in improving COD90 performance.

Introduction

The majority of actions or movements produced in team sport matches require and increase the number of changes of direction (COD) at high speeds to be decisive in performance [1–3] or to have an advantage over opponents during competition [3, 4]. COD is a component

of agility, and it describes a movement wherein no reaction to a stimulus is required and it is previously planned [2, 5]. COD ability is a multifactorial skill which, in its performance, depends on neuromuscular coordination [2], leg muscle strength and power [2, 3, 5] and straight sprinting speed [5–7]. During the COD, it is necessary to execute eccentric force rapidly to decelerate and develop the concentric strength to accelerate in the new direction [1, 2]. In depth, some authors have found that COD correlates with eccentric knee-flexor strength [8] and maximal eccentric lower-body strength [9, 10]. Despite those obvious physical requirements, there is a lack of information regarding the effects of eccentric-overload training on COD performance [2] and its influence on the different factors related to COD performance.

Resistance training produces muscular adaptations at both structural and functional levels [11–14]. Thus, numerous studies have demonstrated that chronic resistance exercise can produce an increase in hypertrophy and strength within the first 4 weeks of training [15, 16]. These adaptations are also shown in several studies that have used training protocols focusing on the concentric (CON) or eccentric (ECC) phases of the movement [12, 17–19]. Nevertheless, the increase in hypertrophy and strength is magnified using protocols in which the actions combine CON-ECC phases [12, 20, 21]. It is known that, in these CON-ECC actions, the capacity to produce force is much higher during the ECC than in the CON phase [22–27]. Therefore, some authors argue that training protocols, in which exercise is overloaded during the ECC phase of the movement, achieve greater strength gains than those in which the load is constant during both CON-ECC phases [27–29]. This has given rise to the concept of "eccentric overload" (EO).

Different training devices, using the inertia of rotating flywheel(s), have been designed to increase the EO during CON-ECC movements [30, 31]. This technology generates resistance by opposing the athlete's effort with the inertial force generated by a lightweight rotating flywheel such that the same inertia must be overcome during each repetition by means of accommodated loading [32]. The training load on this technology can be regulated by increasing the speed of movement or by adding flywheel weights. Several studies have confirmed the efficacy of these training devices for improving hypertrophy [32–36], power [32, 34, 37], countermovement jump (CMJ) performance [9, 37–39], 10-m sprint time [4, 9, 38, 39] and COD [4, 9, 38–40].

In our knowledge, there are few studies that have investigated the effects of unilateral and bilateral strength training on physical performance (i.e., linear sprint, strength, power and COD) in team sport players [41, 42]. Only one study examined the effects of bilateral vs unilateral EO training using inertial devices on physical performance in team sports [4], and thus, the influence of this variable (bilateral vs unilateral EO using inertial devices) on the hypertrophy and COD performance factors is still unclear. Therefore, the aim of the present study was to compare the chronic eccentric-overload training effects of unilateral lunge) vs bilateral (half-squat) using an inertial device on hypertrophy and physical performance (measured as lower limb power, 10-m sprint, CMJ, COD).

Materials and methods

Subjects

Twenty-seven young healthy active males were divided in a unilateral lunge group (UG n = 14; age: 22.8 ± 2.9 years; body mass: 75.3 ± 8.8 kg; height: 177.3 ± 3.7 cm) and a bilateral squat group (BG n = 13; age: 22.6 ± 2.7 years; body mass: 79.5 ± 12.8 kg; height: 164.2 ± 7 cm), based on COD test results. All subjects were team sport players or sports science students and were actively training 3–4 times per week on average. All subjects were fully informed about the

protocol and were required to give written consent in accordance with current national and international laws and regulations governing the use of human subjects (Declaration of Helsinki II) in research. The Institutional Ethics Committee (Pablo de Olavide University, Seville, Spain) approved this study.

Experimental procedures

Magnetic resonance imaging (MRI) and muscle performance tests were measured in the preand post-experimental periods. All participants were familiarized with the test procedures to avoid any learning effects. On the 1st day, a MRI was performed. On the 2nd day, subjects performed the field tests on an indoor futsal court in the following order: CMJ, 10-m sprint, and COD tests. There was a 5-min recovery period between the tests. Seventy-two hours later, lower limb power was assessed using an inertial device in the laboratory. Subjects were asked to abstain from performing any strenuous exercises on the day before each test, and they were also asked to follow a similar diet on the day of the test.

Magnetic resonance imaging. After one hour of supine rest to minimize the influence of a fluid shift on muscle size [43], axial scans were obtained using a 0.2 T MRI (E-scan, Esaote Biomedica, Genoa, Italy); Turbo 3-D T1; time to echo = 16 ms; time to repetition = 40 ms; NSA 1; scan time 3 min 57 s; field of view FOV 180 \cdot 180 mm; matrix 256 \cdot 256 pixels. For each participant, 20 continuous images of the thigh and calf with 6.3-mm-thick slices, with no spacing in between slices, were obtained. The cross-sectional area (CSA) of the *rectus femoris* (RF), the *vastus lateralis* (VL), the *vastus intermedius* (VI), the *vastus medialis* (VM), the *adductor major* (AM), the *lateral gastrocnemius* (LG), the *medial gastrocnemius* (MG), and the *soleus* (S) muscles were analysed in the dominant leg.

Three images were selected for each subject. The first was at the insertion of the AM to analyse the volume of each quadricep muscle. The second was at the junction of the popliteal artery with the fascia of the adductor major to analyse the volume of the AM. The third was coinciding with the section of the greater perimeter of the leg to analyse the volume of LG, MG, P and S. Every image was analysed by manually encircling the individual muscles using OsiriX dicom viewer 2.2.1 software (Osirix Foundation, Ginebra, Suiza). All muscles were encircled separately, but for the analysis, VL and VI appeared together. Subjects also refrained from excessive muscular exercises for 72 h before reporting to the MRI facility.

Field tests. After a 7- to 8-min standardized general warm-up, which included jogging, joint mobility exercises, and 2 sets of 10 repetitions of squats, field tests were carried out. Additionally, a specific warm-up was performed before each test: 1 set of 5 reps (CMJ) and 2 sub-maximal efforts (linear sprint and COD).

Countermovement jump test (CMJ). The CMJ vertical jump test was used to maximize the stretch-shortening cycle activity and assess the explosive strength of the lower limb muscles. The test was performed using an infrared contact platform (Optojump®, Microgate, Bolzano, Italy). During the CMJ, all subjects were instructed to rest their hands on their hips, which was followed by a vertical jump at maximal effort and landing in an upright position keeping their knees extended. The CMJ was performed three times, and it was separated by 45 s of passive recovery. The best jump performance was used for the subsequent statistical analysis.

10-m linear sprint test. Subjects performed a 10-m sprint test, and time was recorded using photoelectric cells (Racetime2, Microgate®, Bolzano, Italy). The front foot was placed 0.5 m behind the first timing gate, and players started voluntarily so the reaction time was eliminated. After a specific warm-up, including the 2 submaximal efforts, two trials were completed. Two minutes of passive recovery occurred between the trials. The best performance trial was used for the subsequent statistical analysis (T-10m).

COD test. COD capacity was measured by four tests. On the one hand, subjects performed two maximum 5 m+5 m straight line sprints with a COD turn of 90°. One of them was done with the dominant leg on the outside (COD90d) during the turn, and the other with the nondominant leg (COD90nd). The turning space was fixed by 4 sticks (height: 1.5 m) which were placed vertically in other to avoid a curvilinear trajectory. On the other, subjects performed two maximum 5 m+5 m straight line sprints with a COD turn of 180°. Subjects performed the COD by touching a line with the dominant leg (COD180d) and with the non-dominant leg (COD180nd). Subjects were encouraged to complete two trials of each test as fast as possible. Two minutes of passive recovery occurred between trials and tests. The time was recorded with the same photoelectric cells (Racetime², Microgate[®], Bolzano, Italy), using two of them in COD90d and COD90nd (placed on the start and finish line) and only one in the start/finish line of the COD180d and COD180nd. As in the 10-m test, the front foot was placed 0.5 m before the first timing gate, and subjects started voluntarily so the reaction time was eliminated. The best time of each test (COD90d, COD90nd, COD180d and COD180nd) was compared with the T-10m to estimate the percentage mean of speed loss due to executing the COD (DEC-COD90d, DEC-COD90nd, DEC-COD180d and DEC-COD180nd), through the formula [(COD-T-10m)/ T-10m) x 100].

Laboratory test. After a 7- to 8-min standardized general warm-up, included jogging, joint mobility exercises and two sub-maximal sets of 8 reps of a half-squat exercise, two lower limb power tests were done. The tests consisted of the assessment of power in the half-squat and lateral lunge exercises using a non-gravity dependent flywheel inertial device (Exxentric kBox, Exxentrix AB, Stockholm, Sweden), allowing subjects to perform maximal CON and ECC actions [44]. Subjects performed 2 monitored sets of all-out 8 repetitions of squats (0.10 kg/m² moment inertia) with both legs (POWER) and a lateral lunge (0.05 kg/m² moment inertia) with dominant (POWERd) and non-dominant leg (POWERnd). The best of the two sets, according to the criteria of higher rotational mean power relative to body weight, was considered for subsequent analysis. To produce EO, subjects were requested to push maximally through the entire range of motion of the CON action and then gently resist during the initial part (first 20°-30°) of the subsequent ECC action before resisting maximally, aiming at bringing the wheels to a stop at ~90° knee angle before initiating the next cycle [45]. There was a 2-min rest in the standing position between each set. As in previous studies with these inertial devices [37], power and velocity during the training session were sampled at 100 Hz using a rotatory encoder (SmartCoach^{TM,} SmartCoach Europe AB, Stockholm, Sweden) and associated software (SmartCoach[®] v.5.2.0.5).

Flywheel resistance exercise training

Participants underwent a 6-week training programme based on bilateral squat (BG) or unilateral lunge (UG) on a flywheel inertial device (Exxentric kBox, Exxentrix AB, Stockholm, Sweden). They performed 2 sessions/week with at least 48 h of rest between the sessions in the same conditions of the test. Each session was structured in a brief standardized warm-up (like that used before in the power test) and 4 sets of 7 repetitions of squat (0.10 kg/m² moment inertia) in BG and a unilateral squat with each leg (0.05 kg/m² moment inertia) in UG. A 3-min recovery period was allowed between sets.

Statistical analysis

Data are presented as mean \pm SD. All data were log-transformed for analysis to reduce bias arising from non-uniformity error and then analysed for practical significance using magnitude-based inferences [46]. The effect size (ES, 90% confidence limit (CL)) in the selected

variables was calculated using the SD. Threshold values for Cohen ES statistics were >0.2 (small), >0.6 (moderate), and >1.2 (large) [46]. The chance that any difference was better/ greater (i.e., greater than the smallest worthwhile change, SWC [0.2 multiplied by the between-subject standard deviation, based on Cohen's d principle, ES]) or similar or worse/ smaller than the other group, was subsequently calculated. Quantitative chances of beneficial/ better or detrimental/poorer effect were assessed qualitatively as follows [46, 47]: <75%, unclear; >75–95%, likely; >95–99%, very likely; and >99%, almost certain. If the chance of having beneficial/better or detrimental/poorer was >75%, the true difference was considered clear (substantial)[48].

Results

The pretreatment MRI analysis showed that the muscle volume of BG was greater in AM (+8.9% [90%CL: -18.21;1.28] moderate ES) and S (+17.1% [90%CL: -25.12;8.31], large ES) than in UG. Likewise, the pretreatment field analysis showed that the DEC-COD90d (+15.4% [90%CL: -6.72;42.62], moderate ES), the DEC-COD90nd (+20.3% [90%CL: -4.4;51.44], moderate ES), the DEC-COD180d (+12.7% [90%CL: -4.85;33.59], moderate ES), and the DEC-COD180nd (+15.1% [90%CL: -3.48;37.34], moderate ES) of UG were greater than in BG.

The pre- and post-analysis of lower-limb MRI and tests of UG and BG are shown in Tables 1 and 2, respectively. Both training groups substantially increased VM and MG muscle volumes (Tables 1 and 2). The UG obtained a greater increased in AM (ES = 0.46 [90%CL: -0.15;1.7], small) and VM (ES = 1.28 [90%CL: 0.15;2.4], large) than BG (Fig 1). The BG substantially increased the muscle volume of the VL + VI (ES = 0.73 [90%CL: -0.21;1.67], moderate) and LG (ES = 0.78 [90%CL: -0.3;1.85], moderate) compared with UG (Fig 1).

In a field test, both training groups substantially increased the CMJ and substantially decreased COD90d and DEC-COD90d without substantial differences between groups (Fig 2). The UG substantially decreased DEC-COD90nd, obtaining substantial differences compared with BG (ES = 0.33 [90%CL: -0.59;1.25], small). Likewise, the UG substantially decreased DEC-COD180d and DEC-COD180Nd, without substantial differences compared with the BG. There was no substantial effect of training in T-10m for each group (Tables 1 and 2), although the UG increased, while the BG did not change T-10m.

In a laboratory test, both groups substantially increased the POWER, POWERd and POWERnd. The increased of POWER in BG (ES = 1.5 [90%CL: -0.42;2.58], moderate) was greater than in UG (Fig 2).

Discussion

The aim of the present study was to compare the effects of chronic eccentric-overload training of unilateral (lateral lunge) vs bilateral (half-squat) using an inertial device in muscle volume, CODs, power, CMJ and 10-m sprint improvements in team sports players. The main findings of this study were as follows. 1. The UG training increased AM and VM muscle volumes sub-stantially more than did BG, while BG increased in VL+VI and LG muscle volume substantially more than did UG. 2. Both groups decreased the time in COD90d, but the UG training provided a better COD 90° transition phase in the non-dominant leg than did BG training; 3. Both types of training increased the power generated in the inertial system during the half-squat and unilateral lunge, but the BG increment was substantially higher than UG in the half-squat test. 4. Both types of training increased the CMJ performance but did not change the time in 10-m sprint test.

Inertial devices have been shown to be very effective in improving muscle size in shortterm strength training periods [49]. In our study, using a flywheel inertial device for only 6

	ONE
--	-----

Variable	Pre	Post	% Diff (CL)	ES (CL)	Chances %	Qualitative magnitude
MRI						
VL+VI	42.16 ± 10.46	43.71 ± 9.75	4.34 (-4.89; 14.46)	0.16 (-0.19; 0.52)	43/53/5	Unclear
RF	7.24 ± 1.86	7.20 ± 2.19	-1.91 (-11,91; 9,22)	-0.07 (-0.47; 0.33)	12/59/29	Unclear
VM	17.14 ± 4.09	19.31 ± 5.14	12.66 (6.35; 18.97)	0,41 (0.21; 0.61)	95/5/0	Very Likely
AM	18.06 ± 3.02	20.08 ± 3.23	11.13 (4.86; 17.77)	0.61 (0.27; 0.95)	98/2/0	Very Likely
MG	15.48 ± 4.17	16.70 ± 4.25	8.22 (3.60; 13.05)	0.27 (0.12; 0.42)	79/21/0	Likely
LG	10.35 ± 2.43	10.43 ± 2.47	0.53 (-9.56; 11.75)	0.02 (-0.41; 0.45)	24/58/19	Unclear
S	28.12 ± 3.84	29.22 ± 4.07	3.81 (-0.42; 8.21)	0,26 (-0.03; 0.56)	65/34/1	Unclear
Field Test						
CMJ (cm)	34.42 ± 4,91	35.98 ± 4,89	4.69 (1,76; 7,70)	0.28 (0.11; 0.45)	78/22/0	Likely
T-10m (s)	1.79 ± 0.19	$1,85 \pm 0.08$	4.14 (-2.91; 11.71)	0.31 (-0.22; 0.84)	64/30/6	Unclear
COD180d (s)	2.70 ± 0.11	2.66 ± 0.12	-1.22 (-2.40; -0.02)	-0.29 (-0.57; -0.01)	0/29/71	Unclear
COD180nd (s)	2.68 ± 0.12	2.64 ± 0.14	-1.53 (-3.35; 0.33)	-0.33 (-0.74; 0.07)	2/26/72	Unclear
COD90d (s)	2.54 ± 0.13	2.44 ± 0.15	-4.04 (-6.74; -1.27)	-0.75 (-1.27; -0.23)	0/4/96	Very likely
COD90nd (s)	2.55 ± 0.16	2.46 ± 0.16	-3.60 (-5.53; -1.65)	-0.54 (-0.83; -0.24)	0/3/97	Very Likely
DEC-COD180d (%)	53.11 ± 23.70	44.08 ± 6.60	-12.93 (-26.56; 3.21)	-0.41 (-0.91; 0.09)	3/21/76	Likely
DEC-COD180nd (%)	52.25 ± 23.71	42.84 ± 7.76	-14.10 (-29.44; 4.57)	-0.43 (-0.99; 0.13)	3/20/76	Likely
DEC-COD90d (%)	44.86 ± 25.67	32.12 ± 7.09	-23.12 (-38.35; -4.14)	-0.60 (-1.11; -0.10)	1/8/91	Likely
DEC-COD90nd (%)	45.41 ± 26.92	33.10 ± 7.53	-21.16 (-34.98; -4.40)	-0.51 (-0.92; -0.10)	0/10/90	Likely
Laboratory Test						
POWER (w/kg)	6.82 ± 1.52	8.14 ± 2.10	18.58 (9.02; 28.96)	0.70 (0.36; 1.05)	99/1/0	Very Likely
POWERd (w/kg)	5.35 ± 1.37	7.70 ± 1.65	44.83 (27.96; 63.92)	1.44 (0.96; 1.92)	100/0/0	Almost Certainly
POWERnd (w/kg)	5.48 ± 1.12	7.46 ± 1.75	35.45 (20.64; 52.07)	1.42 (0.88; 1.97)	100/0/0	Almost Certainly

Table 1. Relative differences and qualitative outcomes in cross-sectional area (CSA), functional test post a unilateral squat training period. Data are mean ± SD.

NOTE: VL+VI = Vastus lateralis and vastus intermedius; RF = rectus femoris; VM = vastus medialis; AM = adductor major; LG = lateral gastrocnemius; MG = medial gastrocnemius; P = Peroneus; S = soleus; CMJ = Couter movement junp; T-10m = time in 10 m linear sprint test; COD180d and COD180nd = time in 5+5 m sprint change of direction of 180° with dominant and non-dominat foot; COD90d and COD90nd = time in 5+5 m sprint change of direction of 90° with dominant and non-dominat foot; DEC-COD = percentage mean speed loss due to execute CODs compared to 10 m sprint time; POWER, POWERd and POWERnd = power relative to bodyweight in squat exercise with both legs, dominant leg and non-dominant leg. CL = confidence limits; ES = effect size; Chances = percentage chance of having better/similar/poorer values.

https://doi.org/10.1371/journal.pone.0193841.t001

weeks, muscle volume increased from 7-8% in MG to 11-12% in MV in both training groups. Only the UG led to an increase in muscle volume of the AM (+ 11%) being substantially greater than that obtained by BG. Likewise, only the BG increased the VL + VI by 10% and LG by 9%, obtaining substantial differences compared with the UG. Previous studies [32, 34–36] using knee extension flywheel inertial devices during similar training periods (5 weeks) have shown similar improvements in the same muscles, but in smaller magnitudes, VM (from 7.9% to 8.6%) and VL + VI (From 5.8% to 7.8%). Furthermore, these studies have also described increases in the rectus femoris volume in both unilateral (From 9.1% to 9.9%, [32, 34, 36]), and bilateral exercises (11.4%, [35]), which did not occur in our study. This finding may be due, at least in part, to the different mechanical demands on the quadricep's musculature. With an open kinetic chain exercise (i.e., using a knee extension flywheel inertial device), in which there is a block of hip movement, there seems to be a more pronounced increase in the RF [32, 34-36]. Nevertheless, in closed kinetic chain exercises (i.e., using a Squat flywheel inertial device), in which there is a hip extension, there seems to be a more pronounced increase in the lateral-medial musculature of the quadriceps [50]. This effect of muscular demand can be appreciated even in the same inertial device by analysing the UG and BG training. In our

	ONE
--	-----

Variable	Pre	Post	% Diff (CL)	ES (CL)	Chances %	Qualitative magnitude
MRI						
VL+VI	43.88 ± 5.67	48.62 ± 8.84	9.98 (3.35; 17.04)	0.62 (0.21; 1.02)	96/4/0	Very Likely
RF	8.17 ± 2.27	8.63 ± 2.14	6.47 (-1.69; 15.31)	0.21 (-0.06; 0.48)	53/46/1	Unclear
VM	18.90 ± 4.40	21.03 ± 4.01	11.26 (2.02; 20.05)	0.46 (0.13; 0.78)	91/9/0	Likely
AM	19.84 ± 3.14	20.56 ± 3.83	3.10 (-1.86; 8.32)	0.18 (-0.11; 0.46)	44/54/2	Unclear
MG	16.59 ± 2.92	17.69 ± 2.85	7.02 (2.17; 12.10)	0.34 (0.11; 0.58)	85/15/0	Likely
LG	9.26 ± 2.06	10.17 ± 2.78	9.09 (-0.04; 19.05)	0.33 (0.00; 0.67)	75/24/1	Likely
S	34.11 ± 5.72	33.47 ± 6.10	-2.07 (-4.75; 0.69)	-0.12 (-0.27; 0.04)	0/83/17	Unclear
Field Test						
CMJ (cm)	34.65 ± 3.78	36.52 ± 4.89	5.12 (1.35; 9.03)	0.42 (0.11; 0.73)	89/11/0	Likely
T-10m (s)	1.85 ± 0.07	1.85 ± 0.09	0.03 (-1.76; 1.85)	0.01 (-0.44; 0.45)	23/56/21	Unclear
COD180d (s)	2.68 ± 0.15	2.66 ± 0.14	-0.66 (-3.54; 2.30)	-0.11 (-0.62; 0.39)	15/47/39	Unclear
COD180nd (s)	2.65 ± 0.15	2.65 ± 0.14	-0.06 (-1.38; 1.28)	-0.01 (-0.23; 0.21)	6/86/8	Unclear
COD90d (s)	2.51 ± 0.11	2.43 ± 0.12	-3.21 (-4.57; -1.84)	-0.70 (-1.00; -0.40)	0/1/99	Very Likely
COD90nd (s)	2.49 ± 0.13	2.45 ± 0.14	-1.63 (-4.12; 0.92)	-0.29 (-0.74; 0.16)	4/33/64	Unclear
DEC-COD180d (%)	44.94 ± 6.93	43.94 ± 6.78	-2.23 (-12.28; 8.97)	-0.14 (-0.81; 0.53)	19/37/44	Unclear
DEC-COD180nd (%)	43.13 ± 6.22	42.95 ± 4.86	0.09 (-6.48; 7.13)	0.01 (-0.40; 0.41)	21/60/19	Unclear
DEC-COD90d (%)	35.77 ± 5.29	31.46 ± 7.24	-13.17 (-20.91; -4.66)	-0.84 (-1.39; -0.28)	0/3/97	Very Likely
DEC-COD90nd (%)	34.53 ± 6.48	32.29 ± 6.34	-6.61 (-16.38; 4.30)	-0.34 (-0.89; 0.21)	5/28/67	Unclear
Laboratory Test						
POWER (w/kg)	$6.61 \pm 1,58$	$9.05 \pm 1,70$	38.63 (29,17; 48,78)	1.15 (0.90; 1.40)	100/0/0	Almost Certainly
POWERd (w/kg)	4.85 ± 1.13	7.35 ± 1.36	53.23 (37.29; 71.01)	1.55 (1.15; 1.94)	100/0/0	Almost Certainly
POWERnd (w/kg)	5.00 ± 1.15	6.95 ± 1.34	40.68 (26.09; 56.95)	1.24 (0.84; 1.64)	100/0/0	Almost Certainly

Table 2. Relative differences and qualitative outcomes in cross-sectional area (CSA) and functional test post a bilateral squat training period. Data are mean ± SD.

NOTE: VL+VI = Vastus lateralis and vastus intermedius; RF = rectus femoris; VM = vastus medialis; AM = adductor major; LG = lateral gastrocnemius; MG = medial gastrocnemius; P = Peroneus; S = soleus; CMJ = Couter movement junp; T-10m = time in 10 m linear sprint test; COD180d and COD180nd = time in 5+5 m sprint change of direction of 180° with dominant and non-dominat foot; COD90d and COD90nd = time in 5+5 m sprint change of direction of 90° with dominant and non-dominat foot; DEC-COD = percentage mean speed loss due to execute CODs compared to 10 m sprint time; POWER, POWERd and POWERnd = power relative to bodyweight in squat exercise with both legs, dominant leg and non-dominant leg. CL = confidence limits; ES = effect size; Chances = percentage chance of having better/similar/poorer values.

https://doi.org/10.1371/journal.pone.0193841.t002

study, there was similar muscular growth in all muscles except in the AM, which increased in UG that performed the lateral lunge exercise. By using 3-D musculoskeletal dynamics simulation models to compare the joint reaction forces, it is known that it is there is a greater demand of the horizontal force in the lateral lunge than in the squat [51]. Furthermore, Fisher and Wallin [52] argued that the use of unilateral strength training can increase the force production of the stabilizing muscles (i.e., hip adductors). These two aspects could partly explain the AM increment in the UG group.

In exploring the relationship between muscle volume and speed, Chelli et al [53] described significant and positive correlations between increased thigh muscle hypertrophy and 5-m sprint speed (r = 0.48) in junior soccer players. However, despite the thigh muscle size increase found in our results, the BG and UG T-10m did not change. Our results are in agreement with those obtained by De Hoyo et al. [38], that reported no improvements in a 10-m sprint after 10 weeks of training using the same bilateral squat exercise in a similar flywheel inertial device. Likewise, Tous-Fajardo et al [39] obtained similar results after 11 weeks of combined unilateral strength training (including a lateral lunge inertial device). Nevertheless, Gonzalo-Skok et al [4] recently compared unidirectional-bilateral vs multidirectional-unilateral exercises using an







inertial device and showed that both types of training-induced substantial enhancements in short distance sprints. The different moment inertia (from 0.05 to 0.1 vs 0.27 kg/m^2), different devices (K-Box vs conical-pulley) and how they were used could explain these differences. In fact, these authors considered that it was very unlikely to obtain EO with a conical-pulley inertial device; moreover, in more stable exercises (i.e., Squat Kbox vs. Squat conical-pulley), a more consistent EO can be developed [4]. Given this controversy caused by both factors, EO





https://doi.org/10.1371/journal.pone.0193841.g002

and stability, together with the fact that a recent meta-analysis has described that an increase in force is significantly higher with the existence of EO during the exercise [49], it could be interesting to study EO during exercise for the improvement of velocity in short distance sprints.

This disparity of results about the effects of UG/BG training on sprint speed is also seen in the COD performance. It has recently been suggested that unilateral and bilateral training may be equally efficacious in improving COD [41]. Nevertheless, some studies have shown substantially greater COD performance improvements after unilateral training using conventional [42, 52] and inertial devices [4]. COD ability is a multifactorial skill in which performance depends on neuromuscular coordination [2], leg muscle strength and power [2, 3, 5] and a straight sprinting speed [5–7]. Regarding the importance of each factor, studies focused on improving COD performance with unilateral/bilateral training do not differentiate if the COD can be improved by increasing all these factors or just by increasing only the straight sprinting speed. We analysed the percentage of speed lost due to execute the COD (DEC-COD) just to differentiate the influence of these factor in the increase of COD performance after training.

In our study, the UG substantially improved both COD90d and COD90nd and all DEC-CODs (90° and 180°), while BG improved only COD90d and DEC-COD90d. The magnitude of these changes in COD90 (ES 0.41 to 0.84) was similar to that recently reported in COD turns of 45° after either unilateral or bilateral training using an inertial device [4]. As in other studies [4, 42], UG always obtained more robust adaptations in all COD tests (i.e., greater mean ES) than BG (Fig 2), but there was only a substantial interaction between groups for DEC-COD90nd. In both our study and in Gonzalo-Skok et al. [42], UG obtained better results in the non-dominant leg than BG. These authors argued there was a difference between dominant and non-dominant vertical ground force that favoured the preferred leg during a bilateral squat. It could be possible that BG might exert more effort with the dominant leg (vs nondominant leg), whilst UG had made the same effort with the dominant and non-dominant leg. As the sprint capacity did not change in our study, COD improvements could depend on good neuromuscular coordination plus muscle strength and power to stop the previous inertia and to accelerate later in the new direction [2]. This effect seems to be reflected in the lower loss of time to stop the previous inertia during the COD90 in both groups (DEC-COD90). It is known that these adaptations depend on the weekly eccentric-overload stimulus elicited by Tous et al [39]. Thus, the fact that BG only improved DEC-COD90d could be partly explained by a different EO stimulation on the dominant and non-dominant leg (as explained before). In this respect, the absence of an electromyographic registry during the training sessions to confirm this suggestion as a limitation of our study.

The CMJ performance was increased from 4.7 to 5.1% (for UG and BG, respectively) without an inter-group effect. This change's magnitude and the lack of inter-group effect were similar to those recently reported by Gonzalo-Skok et al. [4]. However, BG appears to have a higher effect on jumping performance than unilateral training (ES-UG = 0.28; ES-BG = 0.42). Although both exercises improved POWER, POWERd and POWERnd, the BG POWER increment was substantially greater than UG (Fig 2). In this regard, relative strength/power may be essential for COD performance (13). However, this relationship is not observed in our results.

Conclusions

In summary, the results of our study show that 6 weeks of unilateral/bilateral eccentric overload training improved lower limb muscle volume, power, CMJ performance, COD90 and DEC-COD90, without changes in sprint performance in team sport athletes. This finding suggests that EO training provided a better COD turn of the 90° transition phase. Our results also show that UG improved COD90 and DEC-COD90 in the dominant and non-dominant legs, while BG only improved COD90 and DEC-COD90 in the dominant leg. This finding suggests that unilateral training seems to be more effective in improving COD turns of 90° performance.

Author Contributions

Conceptualization: Francisco Javier Núñez.

Data curation: Francisco Javier Núñez, Alfredo Santalla, Irene Carrasquila, Jose Antonio Asian, Jose Ignacio Reina, Luis Jesús Suarez-Arrones.

Formal analysis: Francisco Javier Núñez.

Investigation: Francisco Javier Núñez, Alfredo Santalla, Irene Carrasquila, Jose Antonio Asian, Jose Ignacio Reina, Luis Jesús Suarez-Arrones.

Methodology: Francisco Javier Núñez, Alfredo Santalla, Luis Jesús Suarez-Arrones.

Supervision: Francisco Javier Núñez.

Writing - original draft: Francisco Javier Núñez.

Writing - review & editing: Alfredo Santalla, Luis Jesús Suarez-Arrones.

References

- Ben Abdelkrim N, Chaouachi A, Chamari K, Chtara M, Castagna C. Positional role and competitivelevel differences in elite-level men's basketball players. J Strength Cond Res. 2010; 24(5):1346–55. Epub 2010/04/16. https://doi.org/10.1519/JSC.0b013e3181cf7510 PMID: 20393355.
- Brughelli M, Cronin J, Levin G, Chaouachi A. Understanding change of direction ability in sport: a review of resistance training studies. Sports Med. 2008; 38(12):1045–63. Epub 2008/11/26. <u>https://doi.org/10.2165/00007256-200838120-00007</u> PMID: 19026020.
- Sheppard JM, Young WB. Agility literature review: classifications, training and testing. J Sports Sci. 2006; 24(9):919–32. Epub 2006/08/03. https://doi.org/10.1080/02640410500457109 PMID: 16882626.
- Gonzalo-Skok O, Tous-Fajardo J, Valero-Campo C, Berzosa C, Bataller AV, Arjol-Serrano JL, et al. Eccentric-Overload Training in Team-Sport Functional Performance: Constant Bilateral Vertical Versus Variable Unilateral Multidirectional Movements. Int J Sports Physiol Perform. 2017; 12(7):951–8. Epub 2016/12/15. https://doi.org/10.1123/ijspp.2016-0251 PMID: 27967273.
- Young WB, James R, Montgomery I. Is muscle power related to running speed with changes of direction? J Sports Med Phys Fitness. 2002; 42(3):282–8. Epub 2002/07/03. PMID: 12094116.
- Castillo-Rodriguez A, Fernandez-Garcia JC, Chinchilla-Minguet JL, Carnero EA. Relationship between muscular strength and sprints with changes of direction. J Strength Cond Res. 2012; 26(3):725–32. Epub 2012/02/01. https://doi.org/10.1519/JSC.0b013e31822602db PMID: 22289697.
- Little T, Williams AG. Specificity of acceleration, maximum speed, and agility in professional soccer players. J Strength Cond Res. 2005; 19(1):76–8. Epub 2005/02/12. https://doi.org/10.1519/14253.1 PMID: 15705049.
- Jones P, Bampouras TM, Marrin K. An investigation into the physical determinants of change of direction speed. J Sports Med Phys Fitness. 2009; 49(1):97–104. Epub 2009/02/04. PMID: 19188902.
- de Hoyo M, Sanudo B, Carrasco L, Dominguez-Cobo S, Mateo-Cortes J, Cadenas-Sanchez MM, et al. Effects of Traditional Versus Horizontal Inertial Flywheel Power Training on Common Sport-Related Tasks. J Hum Kinet. 2015; 47:155–67. Epub 2015/11/12. https://doi.org/10.1515/hukin-2015-0071 PMID: 26557200; PubMed Central PMCID: PMCPMC4633251.
- Spiteri T, Nimphius S, Hart NH, Specos C, Sheppard JM, Newton RU. Contribution of strength characteristics to change of direction and agility performance in female basketball athletes. J Strength Cond Res. 2014; 28(9):2415–23. Epub 2014/05/31. <u>https://doi.org/10.1519/JSC.00000000000547</u> PMID: 24875426.
- Aagaard P, Andersen JL, Dyhre-Poulsen P, Leffers AM, Wagner A, Magnusson SP, et al. A mechanism for increased contractile strength of human pennate muscle in response to strength training: changes in

muscle architecture. J Physiol. 2001; 534(Pt. 2):613–23. https://doi.org/10.1111/j.1469-7793.2001.t01-1-00613.x PMID: 11454977; PubMed Central PMCID: PMCPMC2278719.

- Higbie EJ, Cureton KJ, Warren GL 3rd, Prior BM. Effects of concentric and eccentric training on muscle strength, cross-sectional area, and neural activation. J Appl Physiol (1985). 1996; 81(5):2173–81. Epub 1996/11/01. https://doi.org/10.1152/jappl.1996.81.5.2173 PMID: 8941543.
- Moritani T, deVries HA. Neural factors versus hypertrophy in the time course of muscle strength gain. Am J Phys Med. 1979; 58(3):115–30. Epub 1979/06/01. PMID: 453338.
- Narici MV, Roi GS, Landoni L, Minetti AE, Cerretelli P. Changes in force, cross-sectional area and neural activation during strength training and detraining of the human quadriceps. Eur J Appl Physiol Occup Physiol. 1989; 59(4):310–9. Epub 1989/01/01. PMID: 2583179.
- Schuenke MD, Herman JR, Gliders RM, Hagerman FC, Hikida RS, Rana SR, et al. Early-phase muscular adaptations in response to slow-speed versus traditional resistance-training regimens. Eur J Appl Physiol. 2012; 112(10):3585–95. Epub 2012/02/14. <u>https://doi.org/10.1007/s00421-012-2339-3</u> PMID: 22328004.
- Staron RS, Karapondo DL, Kraemer WJ, Fry AC, Gordon SE, Falkel JE, et al. Skeletal muscle adaptations during early phase of heavy-resistance training in men and women. J Appl Physiol (1985). 1994; 76(3):1247–55. Epub 1994/03/01. https://doi.org/10.1152/jappl.1994.76.3.1247 PMID: 8005869.
- Colliander EB, Tesch PA. Effects of eccentric and concentric muscle actions in resistance training. Acta Physiol Scand. 1990; 140(1):31–9. Epub 1990/09/01. <u>https://doi.org/10.1111/j.1748-1716.1990.</u> tb08973.x PMID: 2275403.
- Jones DA, Rutherford OM. Human muscle strength training: the effects of three different regimens and the nature of the resultant changes. J Physiol. 1987; 391:1–11. Epub 1987/10/01. PMID: 3443943; PubMed Central PMCID: PMCPMC1192197.
- Seger JY, Arvidsson B, Thorstensson A. Specific effects of eccentric and concentric training on muscle strength and morphology in humans. Eur J Appl Physiol Occup Physiol. 1998; 79(1):49–57. Epub 1999/ 03/03. https://doi.org/10.1007/s004210050472 PMID: 10052660
- Hather BM, Tesch PA, Buchanan P, Dudley GA. Influence of eccentric actions on skeletal muscle adaptations to resistance training. Acta Physiol Scand. 1991; 143(2):177–85. Epub 1991/10/01. https://doi. org/10.1111/j.1748-1716.1991.tb09219.x PMID: 1835816.
- Hortobagyi T, Hill JP, Houmard JA, Fraser DD, Lambert NJ, Israel RG. Adaptive responses to muscle lengthening and shortening in humans. J Appl Physiol (1985). 1996; 80(3):765–72. Epub 1996/03/01. https://doi.org/10.1152/jappl.1996.80.3.765 PMID: 8964735.
- Crenshaw AG, Karlsson S, Styf J, Backlund T, Friden J. Knee extension torque and intramuscular pressure of the vastus lateralis muscle during eccentric and concentric activities. Eur J Appl Physiol Occup Physiol. 1995; 70(1):13–9. Epub 1995/01/01. PMID: 7729433.
- Enoka RM. Eccentric contractions require unique activation strategies by the nervous system. J Appl Physiol (1985). 1996; 81(6):2339–46. Epub 1996/12/01. <u>https://doi.org/10.1152/jappl.1996.81.6.2339</u> PMID: 9018476.
- 24. Katz B. The relation between force and speed in muscular contraction. J Physiol. 1939; 96(1):45–64. Epub 1939/06/14. PMID: 16995114; PubMed Central PMCID: PMCPMC1393840.
- **25.** Komi PV, Viitasalo JT. Changes in motor unit activity and metabolism in human skeletal muscle during and after repeated eccentric and concentric contractions. Acta Physiol Scand. 1977; 100(2):246–54. Epub 1977/06/01. https://doi.org/10.1111/j.1748-1716.1977.tb05943.x PMID: 888714.
- 26. Tesch PA, Dudley GA, Duvoisin MR, Hather BM, Harris RT. Force and EMG signal patterns during repeated bouts of concentric or eccentric muscle actions. Acta Physiol Scand. 1990; 138(3):263–71. Epub 1990/03/01. https://doi.org/10.1111/j.1748-1716.1990.tb08846.x PMID: 2327260.
- Westing SH, Seger JY. Eccentric and concentric torque-velocity characteristics, torque output comparisons, and gravity effect torque corrections for the quadriceps and hamstring muscles in females. Int J Sports Med. 1989; 10(3):175–80. Epub 1989/06/01. <u>https://doi.org/10.1055/s-2007-1024896</u> PMID: 2777436.
- Doan BK, Newton RU, Marsit JL, Triplett-McBride NT, Koziris LP, Fry AC, et al. Effects of increased eccentric loading on bench press 1RM. J Strength Cond Res. 2002; 16(1):9–13. Epub 2002/02/09. PMID: 11834100.
- Hortobagyi T, Devita P, Money J, Barrier J. Effects of standard and eccentric overload strength training in young women. Med Sci Sports Exerc. 2001; 33(7):1206–12. Epub 2001/07/11. PMID: 11445770.
- Berg HE, Tesch A. A gravity-independent ergometer to be used for resistance training in space. Aviat Space Environ Med. 1994; 65(8):752–6. Epub 1994/08/01. PMID: 7980338.
- Berg HE, Tesch PA. Force and power characteristics of a resistive exercise device for use in space. Acta Astronaut. 1998; 42(1–8):219–30. Epub 2001/09/07. PMID: <u>11541605</u>.

- 32. Norrbrand L, Fluckey JD, Pozzo M, Tesch PA. Resistance training using eccentric overload induces early adaptations in skeletal muscle size. Eur J Appl Physiol. 2008; 102(3):271–81. Epub 2007/10/11. https://doi.org/10.1007/s00421-007-0583-8 PMID: 17926060.
- Alkner BA, Tesch PA. Efficacy of a gravity-independent resistance exercise device as a countermeasure to muscle atrophy during 29-day bed rest. Acta Physiol Scand. 2004; 181(3):345–57. Epub 2004/ 06/16. https://doi.org/10.1111/j.1365-201X.2004.01293.x PMID: 15196095.
- Fernandez-Gonzalo R, Irimia JM, Cusso R, Gustafsson T, Linne A, Tesch PA. Flywheel resistance exercise to maintain muscle oxidative potential during unloading. Aviat Space Environ Med. 2014; 85 (7):694–9. Epub 2014/07/16. PMID: 25022156.
- Seynnes OR, de Boer M, Narici MV. Early skeletal muscle hypertrophy and architectural changes in response to high-intensity resistance training. J Appl Physiol (1985). 2007; 102(1):368–73. Epub 2006/ 10/21. https://doi.org/10.1152/japplphysiol.00789.2006 PMID: 17053104.
- Tesch PA, Ekberg A, Lindquist DM, Trieschmann JT. Muscle hypertrophy following 5-week resistance training using a non-gravity-dependent exercise system. Acta Physiol Scand. 2004; 180(1):89–98. Epub 2004/01/07. https://doi.org/10.1046/j.0001-6772.2003.01225.x PMID: 14706117.
- Gual G, Fort-Vanmeerhaeghe A, Romero-Rodriguez D, Tesch PA. Effects of In-Season Inertial Resistance Training With Eccentric Overload in a Sports Population at Risk for Patellar Tendinopathy. J Strength Cond Res. 2016; 30(7):1834–42. Epub 2015/12/17. https://doi.org/10.1519/JSC. 000000000001286 PMID: 26670989.
- de Hoyo M, Pozzo M, Sanudo B, Carrasco L, Gonzalo-Skok O, Dominguez-Cobo S, et al. Effects of a 10-week in-season eccentric-overload training program on muscle-injury prevention and performance in junior elite soccer players. Int J Sports Physiol Perform. 2015; 10(1):46–52. Epub 2014/06/10. <u>https://</u> doi.org/10.1123/ijspp.2013-0547 PMID: 24910951.
- Tous-Fajardo J, Gonzalo-Skok O, Arjol-Serrano JL, Tesch P. Enhancing Change-of-Direction Speed in Soccer Players by Functional Inertial Eccentric Overload and Vibration Training. Int J Sports Physiol Perform. 2016; 11(1):66–73. Epub 2015/05/06. <u>https://doi.org/10.1123/ijspp.2015-0010</u> PMID: 25942419.
- de Hoyo M, de la Torre A, Pradas F, Sanudo B, Carrasco L, Mateo-Cortes J, et al. Effects of eccentric overload bout on change of direction and performance in soccer players. Int J Sports Med. 2015; 36 (4):308–14. Epub 2014/12/20. https://doi.org/10.1055/s-0034-1395521 PMID: 25525954.
- Speirs DE, Bennett MA, Finn CV, Turner AP. Unilateral vs. Bilateral Squat Training for Strength, Sprints, and Agility in Academy Rugby Players. J Strength Cond Res. 2016; 30(2):386–92. Epub 2015/07/23. https://doi.org/10.1519/JSC.00000000001096 PMID: 26200193.
- Gonzalo-Skok O, Tous-Fajardo J, Suarez-Arrones L, Arjol-Serrano JL, Casajus JA, Mendez-Villanueva A. Single-Leg Power Output and Between-Limbs Imbalances in Team-Sport Players: Unilateral Versus Bilateral Combined Resistance Training. Int J Sports Physiol Perform. 2017; 12(1):106–14. Epub 2016/ 05/04. https://doi.org/10.1123/ijspp.2015-0743 PMID: 27140680.
- Berg HE, Tedner B, Tesch PA. Changes in lower limb muscle cross-sectional area and tissue fluid volume after transition from standing to supine. Acta Physiol Scand. 1993; 148(4):379–85. Epub 1993/08/ 01. https://doi.org/10.1111/j.1748-1716.1993.tb09573.x PMID: 8213193.
- Alkner BA, Tesch PA. Knee extensor and plantar flexor muscle size and function following 90 days of bed rest with or without resistance exercise. Eur J Appl Physiol. 2004; 93(3):294–305. https://doi.org/ 10.1007/s00421-004-1172-8 PMID: 15338217
- Fernandez-Gonzalo R, Lundberg TR, Alvarez-Alvarez L, de Paz JA. Muscle damage responses and adaptations to eccentric-overload resistance exercise in men and women. Eur J Appl Physiol. 2014; 114(5):1075–84. Epub 2014/02/13. https://doi.org/10.1007/s00421-014-2836-7 PMID: 24519446.
- 46. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. Med Sci Sports Exerc. 2009; 41(1):3–13. Epub 2008/12/19. <u>https://doi.org/10.1249/MSS.0b013e31818cb278 PMID: 19092709</u>.
- Batterham AM, Hopkins WG. Making meaningful inferences about magnitudes. Int J Sports Physiol Perform. 2006; 1(1):50–7. Epub 2006/03/01. PMID: 19114737.
- Aughey RJ. Increased high-intensity activity in elite Australian football finals matches. Int J Sports Physiol Perform. 2011; 6(3):367–79. Epub 2011/09/14. PMID: 21911862.
- Nunez Sanchez FJ, Saez de Villarreal E. Does Flywheel Paradigm Training Improve Muscle Volume and Force? A Meta-Analysis. J Strength Cond Res. 2017; 31(11):3177–86. Epub 2017/10/27. <u>https:// doi.org/10.1519/JSC.00000000002095</u> PMID: 29068866.
- Irish SE, Millward AJ, Wride J, Haas BM, Shum GL. The effect of closed-kinetic chain exercises and open-kinetic chain exercise on the muscle activity of vastus medialis oblique and vastus lateralis. J Strength Cond Res. 2010; 24(5):1256–62. Epub 2010/04/14. https://doi.org/10.1519/JSC. 0b013e3181cf749f PMID: 20386128.

- Qian JG, Li Z, Zhang H, Bian R, Zhang S. Effectiveness of Selected Fitness Exercises on Stress of Femoral Neck using Musculoskeletal Dynamics Simulations and Finite Element Model. J Hum Kinet. 2014; 41:59–70. Epub 2014/08/13. https://doi.org/10.2478/hukin-2014-0033 PMID: 25114732; PubMed Central PMCID: PMCPMC4120465.
- 52. Fisher J, Wallin M. Unilateral versus bilateral lower-body resistance and plyometric training for change of direction speed. J Athl Enhancement 3. 2014; 6:2.
- Chelly MS, Fathloun M, Cherif N, Ben Amar M, Tabka Z, Van Praagh E. Effects of a back squat training program on leg power, jump, and sprint performances in junior soccer players. J Strength Cond Res. 2009; 23(8):2241–9. Epub 2009/10/15. https://doi.org/10.1519/JSC.0b013e3181b86c40 PMID: 19826302.